

λ -PRESENTABLE MORPHISMS, INJECTIVITY AND (WEAK) FACTORIZATION SYSTEMS

MICHEL HÉBERT[†]

ABSTRACT. We show that every λ_m -injectivity class (i.e., the class of all the objects injective with respect to some class of λ -presentable morphisms) is a weakly reflective subcategory determined by a functorial weak factorization system cofibrantly generated by a class of λ -presentable morphisms. This was known for small-injectivity classes, and referred to as the “small object argument”. An analogous result is obtained for orthogonality classes and factorization systems, where λ -filtered colimits play the role of the transfinite compositions in the injectivity case. λ -presentable morphisms are also used to organize and clarify some related results (and their proofs), in particular on the existence of enough injectives (resp. pure-injectives).

INTRODUCTION

It is well-known that for every (small) set $\mathcal{N}$ of morphisms in a locally presentable categories $\mathcal{C}$, $(\text{Cof}(\mathcal{N}), \mathcal{N}^\square)$ is a weak factorization system, where the class $\text{Cof}(\mathcal{N})$ of *cofibrations* of $\mathcal{N}$ is the class $\text{ReChPo}(\mathcal{N})$ of all retracts of transfinite compositions of pushouts of the members of $\mathcal{N}$ (complete definitions are recalled below). One may refer to this as the small object argument, originating in homotopy theory in the 60’s (see [Bk, 00]). As a consequence, every small-injectivity class is weakly reflective (and determined by the weak factorization above). This was recently generalized in various directions, for example in [AHRT, 02] and in [Ch, 05].

Our main result (Theorem 1.6 below) generalizes the small object argument as follows. A λ -*presentable morphism* of $\mathcal{C}$ is just a λ -presentable object of the comma category $(A \downarrow \mathcal{C})$ for some object A . We show that for any class $\mathcal{M}$ of λ -presentable morphisms, the class $\mathcal{K} = \mathcal{M}^\Delta$ of all objects which are injective with respect to all members of $\mathcal{M}$ is weakly reflective and determined by a weak factorization system $(\text{ReCh}(\mathcal{N}), \mathcal{N}^\square)$, for some class $\mathcal{N}$ of λ -presentable morphisms. Analogous results are shown for orthogonality classes and their associated reflective subcategories and factorization systems, where λ -filtered colimits play the role of the transfinite compositions in the injectivity case.

Variants of the construction of the transfinite composition used in the theorem will have other applications, for example to give simple proofs of the existence of “sufficiently many” pure subobjects, and of “enough” absolutely pure objects in $\mathcal{C}$.

Date: September 8, 2005.

2000 Mathematics Subject Classification. 18A20, 18A32, 13B22, 13C11, 55U35.

Key words and phrases. finitely presentable morphism, finitely presented morphism, pure morphism, injectivity, locally presentable categories, orthogonality, weak factorization system.

[†]Department of Mathematics, The American University in Cairo, Box 2511, Cairo, Egypt. mhebert@aucegypt.edu.

Much of Section 2 of the paper is devoted to the promotion of the λ -presentable morphisms as a natural concept to organize and simplify various results. For example, we show that FP-injective modules are precisely the modules which are injective with respect to finitely presentable monomorphisms, and also that in a locally presentable category $\mathcal{C}$ with the transferability property, an object A is injective (with respect to all monos) if and only if there exists γ such that A is injective with respect to all γ -presentable monos and all γ -pure monos. We finally note that the existence of enough (pure-) injectives in certain categories rests on the fact that there exists γ such that all (pure) monos in those categories are transfinite composition of γ -presentable (pure) monos.

1. MAIN DEFINITIONS AND RESULTS

For basic definitions and results on locally presentable and accessible categories, we refer the reader to [AR, 94]. For convenience, we recall the following.

If λ is a regular infinite cardinal, an object A of a category $\mathcal{C}$ is *λ -presentable* if the hom-functor $\mathcal{C}(A, -): \mathcal{C} \rightarrow \mathbf{Set}$ preserves λ -filtered colimits. Then, $\mathcal{C}$ is *λ -accessible* if it has all λ -filtered colimits, as well as a (small) set S of λ -presentable objects such that every object in $\mathcal{C}$ is the λ -filtered colimit of a diagram with all its vertices in S . Finally, $\mathcal{C}$ is *locally λ -presentable* if it is λ -accessible and cocomplete (or, equivalently, complete). We write *finitely presentable* for ω -presentable. We recall from [H, 98] the following definition.

Definition 1.1. A morphism $f: A \rightarrow B$ in a category $\mathcal{C}$ is called *λ -presentable* if it is a λ -presentable object of the comma category $(A \downarrow \mathcal{C})$. Given a class $\mathcal{N}$ of morphisms, $\lambda\mathcal{N}$ will denote the class of all λ -presentable morphisms in $\mathcal{N}$.

Note that the comma category $(A \downarrow \mathcal{C})$ is locally λ -presentable for every object A in a locally λ -presentable category $\mathcal{C}$ ([H₂, 04]). This is used in particular to show the following, which we will need throughout the paper:

Theorem 1.2. ([H₂, 04]) *Let $\mathcal{C}$ be a locally λ -presentable category. Then $f: A \rightarrow B$ is λ -presentable if and only if there exists a pushout diagram*

$$\begin{array}{ccc} C & \longrightarrow & D \\ \downarrow & & \downarrow \\ A & \xrightarrow{f} & B \end{array}$$

where C and D are λ -presentable.

Remarks and examples 1.3.

- (1) In finitary varieties, the theorem above amounts to say that a morphism $f: A \rightarrow B$ is finitely presentable when f provides a way to “present B ” by adding less than λ generators and relations to some presentation of A (see [H₁, 04]). Using this, one can see for example that a morphism $f: A \rightarrow B$ in the category **CRng** of commutative rings with unit is finitely presentable if and only if it is (isomorphic to) the canonical homomorphism $A \rightarrow A[x_1, \dots, x_n]/(p_1, \dots, p_m)$ for some polynomials $p_1, \dots, p_m$ over the variables $x_1, \dots, x_n$. In other words, if and only if B is a finitely presented A -algebra, with f as its structure morphism. This coincides with the

- “finitely presented” homomorphisms sometimes met in the Commutative Rings literature (see [P, 03] for example).
- (2) One can also use Theorem 1.2 to show that in the category **Mod- R** of all right R -modules, R a ring, the embedding $A \hookrightarrow B$ of a submodule is finitely presentable if and only if the quotient B/A is a finitely presentable module. Note however that a finitely presentable morphism may not have a finitely presentable kernel.
 - (3) As shown in [H₁, 04], Theorem 1.2 is true “up to a retraction in $(A \downarrow \mathcal{C})$ ” in all λ -accessible categories $\mathcal{C}$ with pushouts. Actually, the proof there can be easily adapted to formulate a version for “multipushouts” (in the sense of Diers’ multicolimits: see [AR, 94], for example). In particular, and more explicitly, this implies that in a locally λ -multipresentable category $\mathcal{C}$, a morphism $f: A \longrightarrow B$ is λ -presentable if and only if it is the retract in $(A \downarrow \mathcal{C})$ of a component of the multipushout of some morphism with λ -presentable domain and codomain. Consequently, in the category **Fld** of fields, the finitely presentable morphisms are just the finitely generated extensions. Much of what follows will hold in this extended context.

Convention 1.4. *For the rest of the paper, unless otherwise specified, $\mathcal{C}$ will be a locally λ -presentable category.*

We will use λ -presentable morphisms to clarify and generalize some constructions and results related to small-injectivity and small-orthogonality classes. The following type of construction has been used at least from the 1960’s. We will use it throughout the paper, so we describe it in detail for easy reference.

1.5. Two factorizations

Given a class $\mathcal{N}$ of morphisms in $\mathcal{C}$, recall that a morphism is a *transfinite composition of morphisms in $\mathcal{N}$* , if it is the canonical morphism $F(0) \longrightarrow \operatorname{colim} F$ of a functor $F: \delta \longrightarrow \mathcal{C}$, where δ is an ordinal (seen as a well-ordered category), $(F(\beta) \longrightarrow F(\beta + 1)) \in \mathcal{N}$ for every $\beta < \delta$, and $F(\beta) = \operatorname{colim}_{\gamma < \beta} F(\gamma)$ for every limit ordinal $\beta < \delta$.


Given $f: A \longrightarrow B$, and $\mathcal{N}$ a class of λ -presentable morphisms which is *stable under pushouts* (i.e., the pushout of a member of $\mathcal{N}$ along any morphism is in $\mathcal{N}$), we will construct two different factorizations $f = f_\lambda f^*$ of f , with f^* a transfinite composition of morphisms in $\mathcal{N}$.

Put $A = A_0$ and $f = f_0$. Given any $\alpha < \lambda$, let $\mathcal{G}_\alpha$ be a skeleton of the category of all factorizations $(h, q): A_\alpha \xrightarrow{h} X \xrightarrow{q} B$ of f_α with $h \in \mathcal{N}$, where a morphism from (h, q) to (h', q') is a morphism $r: X \longrightarrow X'$ in $\mathcal{C}$ such that the following commutes:


$$\begin{array}{ccccc} & & X & & \\ & \nearrow h & \downarrow r & \nwarrow q & \\ A_\alpha & & & & B \\ & \searrow h' & \downarrow & \swarrow q' & \\ & & X' & & \end{array}$$

Let G_α be the set of objects of $\mathcal{G}_\alpha$. We consider the cone $S_\alpha = (h_q \mid (h_q, q) \in G_\alpha)$, and the cone L_α of a representative set of the isomorphic classes of the underlying set of S_α . We emphasize that if (h_q, q) and $(h_{q'}, q')$ are distinct objects in G_α ,


$h_q = h_{q'}$ but $q \neq q'$, then h_q and $h_{q'}$ are distinct in S_α but not in L_α . In the first case (called the *strict* case), we take the colimit (= multiple pushout) of the source S_α , and in the second (*loose*) case, the colimit of L_α . Note that $S_\alpha = L_\alpha$ if f is the unique morphism from A to the terminal object $\mathbf{1}$ of $\mathcal{C}$. In both cases, denote by $A_{\alpha+}$ the colimit object, $g_\alpha = g_{\alpha+}: A_\alpha \rightarrow A_{\alpha+}$ the canonical morphism, and $f_{\alpha+}: A_{\alpha+} \rightarrow B$ the induced morphism:



For $\gamma \geq \alpha$, we define $g_{\alpha\gamma}$ by composition (taking the colimit of the chain when needed). We denote $g_{\alpha\lambda}: A_\alpha \rightarrow A_\lambda$ by k_α , and put $f^* = k_0$.



The factorization $f = f_\lambda f^*$ is the required one. Note that the strict version of the construction is *functorial* in the following sense. Given $u: A \rightarrow A'$, $v: B \rightarrow B'$ and $f': A' \rightarrow B'$ such that $vf = f'u$, and $f'_\lambda f'^*$ the strict factorization of f' , there is a naturally induced $w: A_\lambda \rightarrow A'_\lambda$ such that both squares in the following diagram commute:



We spell out g_0 as a transfinite composition for future reference, as we will consider interesting modifications of the construction later. We well-order the underlying set $S_0^* = \{h_\gamma \mid \gamma < \delta = \delta(S_0)\}$ of S_0 (resp. $L_0^* = \{h_\gamma \mid \gamma < \delta = \delta(L_0)\}$ in the loose case). Then we take the successive pushouts as follows. First put $h_0 = h^1 = p^0$, and for $\gamma > 0$, let p^γ be the pushout of h_γ along h^γ , where $h^{\gamma+1} = p^\gamma h^\gamma$, and for a limit ordinal γ , $h^\gamma: A_0 \rightarrow X^\gamma$ is the canonical morphism

to the colimit X^γ of the chain $(p^\beta \mid \beta < \gamma)$. We illustrate for $\gamma \leq \omega$:

$$(3) \quad \begin{array}{c} \begin{array}{ccccccc} & & & & h^{\omega+1} & & \\ & & & & h^\omega & & \\ & & & & h^2 & & \\ & & & & h^1 & & \\ A_0 & \xrightarrow{h^1} & X_0 = X^1 & \xrightarrow{p^1} & X^2 & \cdots & X^\omega & \xrightarrow{p^\omega} & X^{\omega+1} & \cdots & X^\delta = A_1 \\ & \searrow h_0 & & & & & & & & & \\ & & X_1 & & & & & & & & \\ & \searrow h_1 & & & & & & & & & \\ & & \vdots & & & & & & & & \\ & \searrow h_\omega & & & & & & & & & \\ & & X_\omega & & & & & & & & \end{array} \end{array}$$

Clearly p^γ is in $\mathcal{N}$ for all $\gamma < \delta$.

We will first use 1.5 to obtain a generalization of the small object argument. Before, we need to recall some definitions and notations.

An object K in $\mathcal{C}$ is *injective with respect to* (respectively *orthogonal to*) a morphism $n: A \rightarrow B$ if for all $u: A \rightarrow K$ there exists (resp. a unique) $w: B \rightarrow K$ such that $wn = u$

$$\begin{array}{ccc} A & \xrightarrow{n} & B \\ u \downarrow & \swarrow w & \\ K & & \end{array}$$

We write this as $n \nabla K$ (resp. $n \perp K$). For a class $\mathcal{N}$ of morphisms in $\mathcal{C}$, we define

$$\mathcal{N}^\Delta = \{K \mid n \nabla K \text{ for all } n \in \mathcal{N}\}$$

$$\mathcal{N}^\perp = \{K \mid n \perp K \text{ for all } n \in \mathcal{N}\},$$

and for a class of objects $\mathcal{K}$, we define

$$\mathcal{K}^\nabla = \{n \mid n \nabla K \text{ for all } K \in \mathcal{K}\}$$

$$\mathcal{K}^\top = \{n \mid n \perp K \text{ for all } K \in \mathcal{K}\},$$

The class $\mathcal{N}^\Delta$ is also denoted by $\text{Inj}(\mathcal{N})$ or $\mathcal{N}\text{-Inj}$ in the literature.

Given morphisms p and i , we write $p \sqcap i$ (respectively $p \perp i$) if for every commutative square $vp = iu$, there exists a (resp. unique) morphism w making both triangles commute in the following diagram:

$$\begin{array}{ccc} * & \xrightarrow{p} & * \\ u \downarrow & \swarrow w & \downarrow v \\ * & \xrightarrow{i} & * \end{array}$$

If $\mathcal{N}$ is a class of morphisms, define the classes

$$\sqcap \mathcal{N} = \{p \mid p \sqcap i \text{ for all } i \in \mathcal{N}\}$$

$$\mathcal{N}^\sqcap = \{i \mid p \sqcap i \text{ for all } p \in \mathcal{N}\}$$

$$\mathcal{N}^\uparrow = \{p \mid p \perp i \text{ for all } i \in \mathcal{N}\}$$

$$\mathcal{N}^\downarrow = \{i \mid p \perp i \text{ for all } p \in \mathcal{N}\}.$$

Then a *weak factorization system* (resp. a *factorization system*) in $\mathcal{C}$ is a pair $(\mathcal{E}, \mathcal{M})$ of classes of morphisms such that:

- (1) $\Box \mathcal{M} = \mathcal{E}$ (resp. $\mathcal{M}^\dagger = \mathcal{E}$),
- (2) $\mathcal{E}^\Box = \mathcal{M}$ (resp. $\mathcal{E}^\dagger = \mathcal{M}$), and
- (3) every morphism f in $\mathcal{C}$ has a factorization $f = me$ with $m \in \mathcal{M}$ and $e \in \mathcal{E}$.

(Note: These established notations are difficult to harmonize: the tradition in the weak case has been to represent p and i vertically in the square above, so that what is oriented “up/down” in factorization systems becomes “left/right” in the weak ones.)

A class $\mathcal{K}$ of objects in $\mathcal{C}$ is a λ_m -*injectivity class* (resp. a λ -*injectivity class*) if $\mathcal{K} = \mathcal{N}^\Delta$ for some class $\mathcal{N}$ of λ -presentable morphisms (resp. of morphisms with λ -presentable domains and codomains). λ_m -*orthogonality classes* and λ -*orthogonality classes* are defined similarly, replacing $\mathcal{N}^\Delta$ by $\mathcal{N}^\perp$.

We will use the following notations, given a class $\mathcal{N}$ of morphisms:

- (1) $\text{Ch}(\mathcal{N}) :=$ the class of all transfinite compositions of morphisms in $\mathcal{N}$.
- (2) $\text{Po}(\mathcal{N}) :=$ the class of all pushouts of members of $\mathcal{N}$ (along any morphisms).
- (3) $(A \downarrow \mathcal{N}) :=$ the full subcategory of $(A \downarrow \mathcal{C})$, with $\mathcal{N}$ as its class of objects.
- (4) $\text{Re}(\mathcal{N}) :=$ the class of all retracts in $(A \downarrow \mathcal{C})$ of objects in $(A \downarrow \mathcal{N})$.
- (5) $\text{Cof}(\mathcal{N}) := \text{ReChPo}(\mathcal{N})$ (the class of the *cofibrations* of $\mathcal{N}$).
- (6) $\text{Fc}_\lambda(\mathcal{N}) :=$ the class of all the canonical morphisms $A \longrightarrow \text{colim } U_{\mathcal{D}}$, with $\mathcal{D}$ a λ -filtered subcategory of $(A \downarrow \mathcal{N})$, and $U_{\mathcal{D}}: \mathcal{D} \longrightarrow \mathcal{C}$ its forgetful functor (defined by $U_{\mathcal{D}}(n: A \longrightarrow C) = C$).

The following extends results in [Bk, 00], [HAR, 01] and [Co, 79].

Theorem 1.6. *Let $\mathcal{M}$ be a class of λ -presentable morphisms in $\mathcal{C}$. Let $\mathcal{N} = \lambda\text{-}(\mathcal{M}^{\Delta^\nabla})$, and $\mathcal{N}_1 = \lambda\text{-}(\mathcal{M}^{\top^\perp})$. Then*

- (a) $(\text{ReCh}(\mathcal{N}), \mathcal{N}^\Box)$ *is a (functorial) weak factorization system, and*
- (b) $(\text{Fc}_\lambda(\mathcal{N}_1), \mathcal{N}_1^\perp)$ *is a factorization system.*

Proof. (a) $\mathcal{N}$ is easily seen to be stable under pushouts. Given $f: A \longrightarrow B$, we apply the strict factorization $f = f_\lambda f^*$ in 1.5 to f . Hence $f^* \in \text{Ch}(\mathcal{N})$, and we now show that $f_\lambda \in \mathcal{N}^\Box$.

Consider a commutative square

$$\begin{array}{ccc} X & \xrightarrow{n} & Y \\ u \downarrow & & \downarrow v \\ A_\lambda & \xrightarrow{f_\lambda} & B \end{array}$$

with $n \in \mathcal{N}$. By Theorem 1.2, there exists a pushout square


$$\begin{array}{ccc} C & \xrightarrow{z} & D \\ s \downarrow & \lrcorner & \downarrow t \\ X & \xrightarrow{n} & Y \end{array}$$

with C and D λ -presentable. Then su factorizes through one of the k_α 's (refer to the diagram (1) in 1.5), $su = k_\alpha l$, and we let (l', z') be the pushout of (l, z) , $z': A_\alpha \longrightarrow P$. We prove that $z' \in \mathcal{N}$.


First, z' is λ -presentable, since it is the pushout of z . Then consider $h: A_\alpha \longrightarrow K$, with $K \in \mathcal{M}^\Delta$. Because k_α is a transfinite composition of morphisms in $\mathcal{N}$, there exists $h': A_\lambda \longrightarrow K$ such that $h'k_\alpha = h$. Then there exists $p: Y \longrightarrow K$ with $pn = h'u$ (since $n \in \mathcal{N}$). We have now

$$ptz = pns = h'u s = h'k_\alpha l = hl,$$

so that there is a (unique) $q: P \longrightarrow K$ such that $qz' = h$ (and $ql' = pt$). This shows that $z' \in \mathcal{N}$.



Now, because $vtz = f_\lambda us = f_\lambda k_\alpha l$, the pushout $l'z = z'l$ induces a (unique) morphism $r: P \longrightarrow B$ such that $rz' = f_\lambda k_\alpha = f_\alpha$ (and $rl' = vt$). Hence $A_\alpha \xrightarrow{z'} P \xrightarrow{r} B$ is in G_α , and there exists $z'': P \longrightarrow A_{\alpha+}$ such that $z''z' = g_\alpha$ and $f_\alpha z'' = r$.



Then $us = k_{\alpha+}g_\alpha l = k_{\alpha+}z''z'l = k_{\alpha+}z''l'z$, and the pushout $ns = tz$ induces a (unique) morphism $w: Y \longrightarrow A_\lambda$ such that $wn = u$ and $wt = k_{\alpha+}z''l'$. w is the required diagonal: v is the unique x such that $xn = f_\lambda u$ and $xt = vt$, but we have also $f_\lambda wn = f_\lambda u$ and $f_\lambda wt = f_\lambda k_{\alpha+}z''l' = f_\alpha z''l' = rl' = vt$. We conclude that $f_\lambda w = v$, as needed.

The rest follows a known argument. For $(\text{ReCh}(\mathcal{N}), \mathcal{N}^\square)$ to be a weak factorization system, what remains to be seen is that $\text{ReCh}(\mathcal{N}) = {}^\square(\mathcal{N}^\square)$, since we will then have $(\text{ReCh}(\mathcal{N}))^\square = ({}^\square(\mathcal{N}^\square))^\square = \mathcal{N}^\square$. Its *functoriality* refers to the property with the same name mentioned in 1.5 (diagram (2)).

The inclusion $\text{ReCh}(\mathcal{N}) \subseteq {}^\square(\mathcal{N}^\square)$ is clear, so let $(f: A \longrightarrow B) \in {}^\square(\mathcal{N}^\square)$.

We have seen above that f factorizes as $A \xrightarrow{g} C \xrightarrow{h} B$, with $h \in \mathcal{N}^\square$ and $g \in \text{Ch}(\mathcal{N})$. The commutative square $1_B f = hg$ induces a diagonal d making the

triangles commute in the following diagram

$$\begin{array}{ccc}
 A & \xrightarrow{f} & B \\
 g \downarrow & \searrow d & \downarrow 1_B \\
 C & \xrightarrow{h} & B
 \end{array}$$

so that $hd = 1_f$ in $(A \downarrow \mathcal{C})$. Hence $hg = f$ is in $\text{ReCh}(\mathcal{N})$.

(b) Given $f: A \rightarrow B$, let $\mathcal{D}$ be the subcategory of $(A \downarrow \mathcal{N}_1)$ with its objects the morphisms appearing in the cone S_0 (refer to 1.5, with $\mathcal{N} = \mathcal{N}_1$), and its morphisms those of the category $\mathcal{G}_0$. One can verify that $\mathcal{D}$ is λ -filtered in $(A \downarrow \mathcal{C})$. Taking the colimit of the forgetful functor $U_{\mathcal{D}}: \mathcal{D} \rightarrow \mathcal{C}$, the canonical morphisms $A \xrightarrow{h} \text{colim } U_{\mathcal{D}} \xrightarrow{g} B$ gives the required factorization in one step. The verification that the induced morphism g is in $\mathcal{N}^\perp$ is similar to the one in the weak case, as well as the rest of the proof that $(\text{Fc}_\lambda(\mathcal{N}_1), \mathcal{N}_1^\perp)$ is a factorization system. $\square$

Remarks and examples 1.7.

- (1) Let $\mathcal{M}$ be any (small) set of morphisms. We can assume that members of $\mathcal{M}$ have λ -presentable domains and codomains. Then Proposition 1.3 of [Bk, 00] says that $(\text{Cof}(\mathcal{M}), \mathcal{M}^\square)$ is a weak factorization system. This amounts to replace $\lambda(\mathcal{M}^{\Delta^\nabla})$ by $\text{Po}(\mathcal{M})$, since $(\text{Po}(\mathcal{M}))^\square = \mathcal{M}^\square$. Similarly, Corollary 3.3.4 of [Co, 79] shows that $\lambda(\mathcal{M}^{\top\perp})$ can also be replaced by $\text{Po}(\mathcal{M})$ in this case (so that $(\text{Fc}_\lambda \text{Po}(\mathcal{M}), \mathcal{M}^\perp)$ is a factorization system), provided $\mathcal{M}$ admits a λ -strong left calculus of fractions (see II.2 in [HAR, 01]; note that the set of morphisms in $\mathcal{M}^{\top\perp}$ which have λ -presentable domains and codomains does admit a λ -strong left calculus of fractions, as well as $\lambda(\mathcal{N}^{\top\perp})$ for any class $\mathcal{N}$ of λ -presentable morphisms.)

Both facts are proved following the same line than in 1.6, but the proof is simpler in this case.

- (2) Of course we have, in part (a) of the theorem, that $\text{ReCh}(\mathcal{N}) = \text{Cof}(\mathcal{N})$ (since $\text{Po}(\mathcal{N}) = \mathcal{N}$), so that the weak factorization system is *cofibrantly generated* by a class of λ -presentable morphisms. An interesting problem would be to find conditions under which it is cofibrantly generated by some *set*. More generally, one would like to be able to describe $\text{ReCh}(\lambda(\mathcal{M}^{\Delta^\nabla}))$ in a more constructive way, from the elements of $\mathcal{M}$.

A subcategory $\mathcal{K}$ of $\mathcal{C}$ is *weakly reflective* if for every $A \in \mathcal{C}$, there exists $r_A: A \rightarrow A^*$ in $\mathcal{K}^\nabla$ with $A^* \in \mathcal{K}$. If $\mathcal{K}$ is also closed under retracts, we say it is *almost reflective*. In locally λ -presentable categories, we know that:

- (1) ([AR, 94]) Reflective (resp. almost reflective) subcategories are orthogonality (resp. injectivity) classes.
- (2) ([H₁, 04], [H₂, 04]) λ_m -orthogonality (λ_m -injectivity) classes are reflective (almost reflective).

In (2), the fact that a λ_m -injectivity class is almost reflective follows from it being closed under products, because its inclusion in $\mathcal{C}$ satisfies the Solution Set Condition (by [H₂, 04], Lemma 4.2).

If $\mathcal{M}$ is a family of λ -presentable morphisms, then it is easily seen that $(\lambda(\mathcal{M}^{\Delta^\nabla}))^\Delta = \mathcal{M}^\Delta$ and $(\lambda(\mathcal{M}^{\top\perp}))^\top = \mathcal{M}^\top$. Since a morphism $B \rightarrow \mathbf{1}$ to the

terminal object is in $\mathcal{M}^\perp$ iff $B \in \mathcal{M}^\top$ (resp. is in $\mathcal{M}^\square$ iff $B \in \mathcal{M}^\Delta$), the (weak) factorization system of Theorem 1.6 determines a (almost) reflective subcategory $\mathcal{K} = \mathcal{M}^\perp$ (resp. $\mathcal{K} = \mathcal{M}^\nabla$), with (weak) reflectors $r_A: A \longrightarrow R(A)$ from the appropriate factorization $A \xrightarrow{r_A} R(A) \longrightarrow \mathbf{1}$. Concerning the weak case, as mentioned in 1.5, the strict and the loose factorizations of $A \longrightarrow \mathbf{1}$ are the same, and R and r define respectively a functor $R: \mathcal{C} \longrightarrow \mathcal{C}$ and a natural transformation $r: 1_{\mathcal{C}} \longrightarrow R$. Note that, in contrast with the reflective case, weak reflectors are generally not functorial in this sense (see [T, 01] for more on this).

It is well-known that every reflective subcategory of a locally presentable category is determined in this way by some factorization system (see [CHK, 85]). Whether this holds for almost reflective subcategories and weak factorization system appears to be an open problem. However, we conclude from the above:

Corollary 1.8. *Let α be a regular cardinal, $\alpha \geq \lambda$. Then every almost reflective subcategory $\mathcal{K}$ of $\mathcal{C}$ which is an α_m -injectivity class is induced by a functorial weak factorization system $(\text{Cof}(\mathcal{N}), \mathcal{N}^\square)$ for some class $\mathcal{N}$ of α -presentable morphisms (i.e., $\mathcal{K} = \mathcal{N}^\Delta$). (In particular, $\mathcal{K}$ is functorially almost reflective).*

As for the reflective case, note that the proof of Theorem 1.6(b) also gives a construction of the reflectors, extending [HAR, 01], II.3.

Proposition 1.9. *Let $\mathcal{M}$ be a class of λ -presentable morphisms, $\mathcal{K} = \mathcal{M}^\Delta$. Then the following are equivalent:*

- (i) $\mathcal{K}$ is a λ -injectivity class,
- (ii) $\mathcal{K}$ is closed under λ -filtered colimits,
- (iii) every $m \in \mathcal{M}$ is the pushout of some morphism in $\mathcal{K}^\nabla$ with λ -presentable domains and codomains.
- (iv) $\mathcal{K}$ is determined by some (functorial) weak factorization system $(\text{Cof}(\mathcal{N}), \mathcal{N}^\square)$, for some set $\mathcal{N}$ of morphisms with λ -presentable domains and codomains.

The same is true for $\mathcal{K} = \mathcal{M}^\perp$, if one replaces $\mathcal{K}^\nabla$ by $\mathcal{K}^\top$, injectivity by orthogonality, and the weak factorization system $(\text{Cof}(\mathcal{N}), \mathcal{N}^\square)$ by the factorization system $(\text{Fc}_\lambda\text{Po}(\mathcal{N}), \mathcal{N}^\perp)$.

Proof. (i) $\Rightarrow$ (ii) is straightforward. (ii) $\Rightarrow$ (iii) is Lemma 3.6 of [H₁, 04], together with Theorem 1.2 above. (iii) $\Rightarrow$ (iv): let $\mathcal{N}$ be the set of all morphisms in $\mathcal{K}^\nabla$ with λ -presentable domains and codomains, which have some pushout in $\mathcal{M}$. Then $\mathcal{N}^\Delta = \mathcal{M}^\Delta = \mathcal{K}$. That $(\text{Cof}(\mathcal{N}), \mathcal{N}^\square)$ is a weak factorization system is [Bk, 00]. (iv) $\Rightarrow$ (i) is trivial.

The orthogonality case is completely analogous (the argument in the proof of Lemma 3.6 of [H₁, 04] works the same). $\square$

2. EXAMPLES AND APPLICATIONS

Example 2.1. $\mathcal{M} = \lambda\text{-Mor}$.

Let $\mathcal{M}$ be the class $\lambda\text{-Mor}$ of all λ -presentable morphisms in a locally λ -presentable category $\mathcal{C}$. Obviously $\lambda(\mathcal{M}^{\Delta\nabla}) = \mathcal{M}$. Given $f: A \longrightarrow B$, we compare the two factorizations $f = f_\lambda f^*$ in 1.5.

In the strict case, we have $f_\lambda \in \mathcal{M}^\square$ (Theorem 1.6), which mean that for every commutative square

$$\begin{array}{ccc} C & \xrightarrow{z} & D \\ s \downarrow & & \downarrow t \\ A_\lambda & \xrightarrow{f_\lambda} & B \end{array}$$

where z is λ -presentable, there exists $d: D \rightarrow A_\lambda$ such that $dz = s$ and $f_\lambda d = t$ (This will actually force f_λ to be an isomorphism: see below). Looking at the proof of 1.6, one sees that in the case of the loose factorization, everything works the same, except that the found diagonal d will only be guaranteed to satisfy $dz = s$. This means precisely that f_λ is a λ -pure mono (by Theorem 1.2, this is equivalent to the definition in [AR, 94]: see also [H, 98]).

The loose factorization does not lead to any weak factorization system, but what is interesting in this case is the control that one keeps on the presentability of the A_α 's, because each L_α is a cone made of (essentially) distinct λ -presentable morphisms: using Proposition 2.3.11 of [MP, 89] and Theorem 1.2, we find easily a cardinal γ , depending on $\mathcal{C}$ only, such that if A is β -presentable for some $\beta > (\gamma^{<\gamma})^+$, then A_γ is also β -presentable. Regarding $\mathcal{C}$ as a locally γ -presentable category, and applying the loose factorization to $f: A \rightarrow B$, then A_γ is β -presentable and f_γ is a γ -pure mono, hence λ -pure. This simplifies the proof of Theorem 2.33 of [AR, 94] for the existence of “sufficiently many” λ -pure subobjects in $\mathcal{C}$. Note however that pushouts are needed in our case, while Theorem 2.33 of [AR, 94] applies to all accessible categories $\mathcal{C}$.

Back to the strict case, consider our $f_\lambda \in \mathcal{M}^\square$. As any morphism with domain A_λ , it is the colimit of a λ -filtered diagram in $(A_\lambda \downarrow \mathcal{C})$ made of λ -presentable morphisms (see [H₂, 04], Proposition 2.6). From straightforward diagram chasing, we find easily a right inverse to f_λ ; since it is (pure) mono, it is an isomorphism.

Hence the weak factorization system $(\text{ReCh}(\mathcal{M}), \mathcal{M}^\square)$ of the theorem is just the trivial factorization system (Mor, Iso) , and $\mathcal{M}^\Delta$ is $\mathcal{C}$ itself. Note that this implies in particular that every morphism in $\mathcal{C}$ is a transfinite composition of λ -presentable morphisms. We will see in 2.5 that the same property for (pure) monomorphisms is uncommon: it will be used crucially to deduce the existence of “enough (pure-) injectives” in some categories.

Example 2.2. Integral closure.

In [Be, 67], A. Besserre constructs an “integral closure” of rings, which can be described more easily by the construction 1.5. In $\mathcal{C} = \mathbf{CRng}$, let $\mathcal{M}$ be the class of all morphisms of the form $A \rightarrow A[x]/(p)$, with p a monic polynomial. Such a morphism is easily seen to be a finitely presentable mono, actually the pushout of the canonical homomorphism from the free ring on the set $\{a_0, \dots, a_{n-1}\}$ of the coefficients of p , to the ring freely presented by the set of generators $\{a_0, \dots, a_{n-1}, x\}$ and the relation $p(x) = 0$. By Proposition 1.9, $\mathcal{M}^\Delta$ is an ω -injectivity class, and the construction in 1.5 shows that the associated weak reflectors are monos (they actually have several other interesting properties: see [P, 03], where the members of $\mathcal{M}^\Delta$ are called the *absolutely integrally closed rings*).

Example 2.3. $\mathcal{M} = \lambda\text{-Mono}$.

Let $\mathcal{M}$ be the class λ -Mono of all λ -presentable monomorphisms. For example, in the category $\mathbf{Mod}\text{-}R$ of all right R -modules, with $\lambda = \omega$, $\mathcal{M}^\Delta$ is the class of the *FP-injective* modules: this can be seen by comparing Theorem 5.39 in [NY, 03] with 1.3 (2) above. (The FP-injective modules are also called *absolutely pure* modules, but this terminology is misleading in the present context: see below). More generally, recall that an object in a locally λ -presentable category is called *λ -injective* ([Fa, 75]) if it is injective with respect to all monomorphisms with λ -generated domain and λ -presentable codomain (where A is *λ -generated* if the hom-functor $\mathcal{C}(A, -): \mathcal{C} \rightarrow \mathbf{Set}$ preserves colimits of λ -filtered diagrams of monomorphisms). We have:

Proposition. *Let $\lambda_g\text{-Mono}$ be the class of all monomorphisms with λ -generated domain and λ -presentable codomain in $\mathcal{C}$. Then*

$$\lambda_g\text{-Mono} \subseteq \lambda\text{-Mono} \subseteq \text{Po}(\lambda_g\text{-Mono}).$$

In particular, an object A is λ -injective iff it is injective with respect to all λ -presentable monos.

Proof. From [AR, 94], the λ -generated objects are precisely the strong quotients of the λ -presentable objects. Also, it is easy to check that for any diagram $X \xrightarrow{e} Z \xrightarrow{g} Y$ with e a strong epi, g is λ -presentable if ge is. The first inclusion then follows.

Let $f: A \rightarrow B$ be a λ -presentable monomorphism. By Theorem 1.2, there exists a pushout

$$\begin{array}{ccc} C & \xrightarrow{g} & D \\ \downarrow u & & \downarrow v \\ A & \xrightarrow{f} & B \end{array}$$

with C and D λ -presentable. Take the (StrongEpi, Mono) factorization $C \xrightarrow{e} E \xrightarrow{m} D$ of g . Since f is mono, there exists a unique $d: E \rightarrow A$ making everything commute in the obtained diagram. E is λ -generated, and one verifies easily that f is the pushout of m along d . The last statement follows immediately. $\square$

Hence $(\lambda\text{-Mono})^\Delta$ is the class of all λ -injective objects. This means that it is actually a small-injectivity class, since there is only (essentially) a set of λ -generated objects in $\mathcal{C}$. (Actually, one cannot expect to prove that a given λ_m -injectivity class is not a small-injectivity class without using some large-cardinal principle, as this would violates the Vopenka's principle: see [AR, 94], Theorem 6.27). From the proposition above, we have $(\lambda_g\text{-Mono})^\square = (\lambda\text{-Mono})^\square$, $\text{Po}(\lambda_g\text{-Mono}) = \text{Po}(\lambda\text{-Mono})$, and hence

$$(\text{Cof}(\lambda_g\text{-Mono}), (\lambda_g\text{-Mono})^\square) = (\text{Cof}(\lambda\text{-Mono}), (\lambda\text{-Mono})^\square).$$

Note however that $(\lambda\text{-Mono})^\Delta$ is not necessarily a λ -injectivity class. For example, we will see below that the class of all FP-injectives R -modules is an ω -injectivity class (in $\mathbf{Mod}\text{-}R$) if and only if R is a coherent ring.

2.3.1. Enough λ -injectives.

We say that $\mathcal{C}$ *has enough λ -injectives* if each $A \in \mathcal{C}$ is the subobject of some λ -injective. This is easily seen to be equivalent to the weak reflectors $r_A: A \rightarrow A_\lambda$ determined by the above weak factorization system to be monos. We observe that if $\mathcal{C}$ has enough λ -injectives, then $\lambda\text{-}((\lambda\text{-Mono})^{\Delta\nabla}) = \lambda\text{-Mono}$, so that the weak factorization system provided by Theorem 1.6 is just $(\text{Cof}(\lambda\text{-Mono}), (\lambda\text{-Mono})^\square)$ above: indeed, $\lambda\text{-}((\lambda\text{-Mono})^{\Delta\nabla}) \supseteq \lambda\text{-Mono}$ is clear, and if $g: A \rightarrow C$ is in $\lambda\text{-}((\lambda\text{-Mono})^{\Delta\nabla})$, then it must be mono because $r_A: A \rightarrow A_\lambda$ factorizes through it.

Now, assume that every transfinite composition of monos in $\mathcal{C}$ is mono (as in all locally *finitely* presentable categories, for example), and that $\mathcal{C}$ has the *transferability property*, i.e., the class Mono of all monomorphisms is stable under pushouts (one easily shows that this is equivalent to $\lambda\text{-Mono}$ being stable under pushouts). Then, as shown in [Fa, 75], $\mathcal{C}$ has (functorially) enough λ -injectives: the left-hand part f^* of the $(\text{Cof}(\lambda\text{-Mono}), (\lambda\text{-Mono})^\square)$ -factorization of any morphism f (= the strict factorization in 1.5 with $\mathcal{N} = \lambda\text{-Mono}$) is clearly mono in this case. Actually, since $\mathcal{C}$ is locally γ -presentable for any regular $\gamma > \lambda$, one sees easily that for each $A \in \mathcal{C}$, we have a transfinite chain

$$A \rightarrow A_\lambda \rightarrow A_\gamma \rightarrow \dots,$$

where the composition $A \rightarrow A_\gamma$ is the weak reflector associated with $(\text{Cof}(\gamma\text{-Mono}), (\gamma\text{-Mono})^\square)$. Note that if each such chain *weakly stabilizes*, i.e., there exists γ such that $A_\gamma \rightarrow A_\zeta$ is a split mono for all $\zeta > \gamma$, then $\mathcal{C}$ has *enough injectives*: this means that each A is the subobject of some *injective* (= a member of $(\text{Mono})^\Delta$).

Of course, if $\mathcal{C}$ does not have these nice properties, we can always apply Theorem 1.6 to $\mathcal{N} = \lambda\text{-}(\mathcal{M}^{\Delta\nabla})$, but the associated weak reflectors might then not be monos.

Assume that $\mathcal{C}$ has enough λ -injectives. Now, if $(\lambda\text{-Mono})^\Delta$ is a λ -injectivity class, Proposition 1.9 implies in particular that every monomorphism with λ -generated domain and λ -presentable codomain is the pushout of some monomorphism with λ -presentable domain and codomain. One can deduce from that that every λ -generated subobject of a λ -presentable object in $\mathcal{C}$ is λ -presentable. This conclusion is also reached in Theorem 4-15 of [Fa, 75], where such categories are called locally λ -coherent. However the category $\mathbf{Mod}\text{-}R$ has the transferability property, and hence has enough ω -injectives, but it is locally finitely coherent if only if R is a coherent ring (see again [Fa, 75]).

2.3.2. Absolutely λ -pure objects.

An object A is called *absolutely λ -pure* if every monomorphism from A is λ -pure. A λ -injective object is always absolutely λ -pure (use the fact that the right-end morphism in the $(\text{StrongEpi}, \text{Mono})$ -factorization of a λ -presentable morphism is also λ -presentable). We note a simple but interesting consequence (in what follows, A is *λ -pure-injective* if it is injective with respect to all λ -pure monos):

Corollary. *Suppose that $\mathcal{C}$ satisfies the transferability property. Then an object in $\mathcal{C}$ is injective if and only if it is λ -injective and λ -pure-injective.*

Proof. For the non-trivial direction, given $g: A \rightarrow M$ with M λ -injective, and a mono $f: A \rightarrow B$, the pushout f' of f along g is mono, hence λ -pure (since M is absolutely λ -pure). If M is also λ -pure-injective, then f' is a split mono, and the result follows. $\square$

An absolutely λ -pure object is not necessarily λ -injective. More precisely, we know from [Fa, 75] that if transfinite compositions of monos in $\mathcal{C}$ are mono, then the following are equivalent: (i) $\mathcal{C}$ has enough λ -injectives; (ii) the class λ -Mono is stable under pushouts; (iii) $\mathcal{C}$ has the transferability property; (iv) the absolutely λ -pure objects are λ -injective. Nevertheless, we have:

Proposition. *Assume that transfinite compositions of monos in $\mathcal{C}$ are mono. Then $\mathcal{C}$ has enough absolutely λ -pure objects.*

Proof. The conclusion means that for each A , there exists a monomorphism $A \rightarrow A^*$ with A^* absolutely λ -pure. Our proof is strongly inspired by the one of the existence of enough existentially closed objects in [Fa, 75] (Theorem 6-3). However the use of presentable morphisms will simplify it much. We apply a modified version of 1.5 (with $\mathcal{N} = \lambda$ -Mono and $f: A \rightarrow \mathbf{1}$), which seems to be of general interest. To accommodate for the fact that $\mathcal{N}$ is not stable under pushouts, we simply apply the rule: when taking the successive pushouts from the well-ordered L_α , just discard the unpleasant results.

More precisely, and referring to the diagram (3) in 1.5, if the first pushout p^1 (of h_1 along h^1) is not in $\mathcal{N}$ (i.e., is not a mono), replace it by the identity on X^1 (and hence $h^2 = h^1$). Similarly for the pushout of h_2 along h^2 , etc. Then $f^*: A_0 \rightarrow A_\lambda$ is a mono, and we show that A_λ is absolutely λ -pure as follows: given a commutative square

$$\begin{array}{ccc} C & \xrightarrow{z} & D \\ u \downarrow & & \downarrow v \\ A_\lambda & \xrightarrow{g} & B \end{array}$$

with g mono, and C and D λ -presentable, we must find a diagonal from D to A_λ making the upper triangle commute (here we use the definition of purity in [AR, 94]). If $C \xrightarrow{m} E \xrightarrow{e} D$ is the (StongEpi, Mono) factorization of z , then there exists a unique $s: E \rightarrow A_\lambda$ making everything commute. E being λ -generated, s must factorize through one of the k_α 's (see diagram (1) in 1.5), $s = k_\alpha l$, and we let (l', z') be the pushout of (l, z) , $z': A_\alpha \rightarrow P$. Then z' is mono because gk_α factorizes through it. Also, the pushout of z' along any $h^\beta: A_\alpha \rightarrow X^\beta$ is a mono, since $X^\beta \rightarrow B$ must factorize through it. This means that z' was not discarded, and it implies that $A_\alpha \rightarrow A_{\alpha+}$ must factorize through it. The morphism $D \xrightarrow{l'} P \rightarrow A_\lambda$ is then the required diagonal. $\square$

If $\mathcal{C}$ has the amalgamation property, i.e., the pushout of a mono along a mono is a mono (the category of groups is an example), the construction in the proof is just the (unmodified) one in 1.5. Note that the absolutely λ -pure objects may still not form a weakly reflective subcategory of $\mathcal{C}$ in this case, but $f^*: A \rightarrow A_\lambda$, where $f_\lambda f^*$ is the factorization of $A \rightarrow \mathbf{1}$, has the injectivity property with respect to monos: for every *mono* $g: A \rightarrow C$ with C absolutely λ -pure, there exists h with $hf^* = g$.

In **CRng** (which does not even have the amalgamation property), a finitely presentable morphism $f: A \rightarrow B = A[x_1, \dots, x_n]/(p_1, \dots, p_m)$ is a mono if and only if the set $\{p_1, \dots, p_m\}$ of polynomials is “consistent” over A , in the sense that the p_i 's have a common root in *some* extension of A . Hence the construction above

is a kind of “algebraic closure”. But the term “algebraically closed” is confusing here: although a field can be seen to be algebraically closed in the usual sense if and only if it is absolutely pure in the category **CRng** (using the Nullstellensatz, see [Fa, 75]), a field K can be algebraically closed in an extension field L without the embedding being pure. This simply means that there exists a polynomial on *several* variables with coefficients in K which have a solution in L but not in K (see [Po, 79] for an example).

Example 2.4. $\mathcal{M}_\delta = \delta\text{-}\omega\text{Pure}$.

For simplicity, we assume that $\mathcal{C}$ is locally *finitely* presentable. Given a regular infinite cardinal δ , we let $\mathcal{M}_\delta = \delta\text{-}\omega\text{Pure}$ be the class of all δ -presentable monos which are $(\omega\text{-})$ pure (NOT to be confused with the δ -pure monos above). We show that there exists γ (depending on $\mathcal{C}$ only) such that $\mathcal{C}$ has “enough” $(\delta\text{-}\omega\text{Pure})$ -injectives for every $\delta \geq \gamma$, i.e., such that for each A there exists a pure mono $A \longrightarrow A^*$ with A^* in $(\mathcal{M}_\delta)^\Delta$.

The proof of Theorem 2.4 of [BR, 05] (referring back to [AR, 94]), states that there exists $\gamma \geq \lambda$ such that for any $\delta \geq \gamma$, each object C of $\mathcal{C}$ is a δ -filtered colimit $(u_i: C_i \longrightarrow C)$ with the C_i ’s δ -presentable and the u_i ’s pure monos. Regarding $\mathcal{C}$ as locally δ -presentable, it follows from the proof of Theorem 1.2 (see Lemma 2.5 in [H₂, 04]), that for every δ -presentable pure mono f , there exists a pushout diagram

$$\begin{array}{ccc} C & \xrightarrow{z} & D \\ u \downarrow & & \downarrow v \\ A & \xrightarrow{f} & B \end{array}$$

where C and D are δ -presentable and u and v are pure monos. This implies that vz is pure, and hence z too. Since pure monos are stable under pushouts (see [AHT, 96]), we have $\delta\text{-}\omega\text{Pure} = \text{Po}(\mathcal{N})$ for the set $\mathcal{N}$ of the pure monos with δ -presentable domains and codomains. In addition, the induced weak reflectors are pure monos, by their construction (using [H₂, 04], 2.12(9)).

Note also that, just as in 2.3.1, it follows that the weak factorization system provided by Theorem 1.6 is just

$$(\text{ReCh}(\delta\text{-}\omega\text{Pure}), (\delta\text{-}\omega\text{Pure})^\square) = (\text{Cof}(\mathcal{N}), \mathcal{N}^\square),$$

and that for each A we have a transfinite chain

$$A \longrightarrow A_\delta \longrightarrow A_\gamma \longrightarrow \dots,$$

($\gamma > \delta$) where the composition $A \longrightarrow A_\gamma$ is the (pure mono) weak reflector determined by the corresponding weak factorization system. If each such chain weakly stabilizes, then $\mathcal{C}$ has *enough pure-injectives*: each A is a pure subobject of a pure-injective.

Example 2.5. Categories with enough (pure-) injectives.

Again we assume that $\mathcal{C}$ is locally finitely presentable. We follow the line of the proof of Theorem 2.4 in [BR, 05], to give parallel proofs of the facts that the existence of *effective union* of subobjects (resp. pure subobjects) implies the existence of enough (resp. pure-) injectives ([BR, 05] deals with the pure-injectivity case).

Nothing will be really new here, but the use of γ -presentable morphisms allows to see interesting connections, and will shorten the proof in [BR, 05].

Suppose that there exists γ such that $\mathcal{C}$ satisfies the two conditions:

- (1) $\mathcal{C}$ has enough γ -injectives (resp. enough $(\gamma\text{-}\omega\text{Pure})$ -injectives), and
- (2) every mono (resp. pure mono) in $\mathcal{C}$ is the transfinite composition of γ -presentable monos (resp. γ -presentable pure monos).

Then it is obvious from the definitions that $\mathcal{C}$ has enough injectives (resp. pure-injectives) – actually functorially so, since $(\gamma\text{-Mono})^\Delta$ and $(\gamma\text{-}\omega\text{Pure})^\Delta$ are γ_m -injectivity classes.

From 2.4, $\mathcal{C}$ always satisfy (1) for the pure-injectivity case, and from 2.3.1, it satisfies (1) for the injectivity case if (and only if) it satisfies the transferability property. We don't know if condition (2) is necessary for having enough injectives or pure-injectives (or at least functorially so), but it seems to be an interesting property to consider in its own right (see for example 2.1 and 1.3 above).

We now assume that subobjects (resp. pure subobjects) in $\mathcal{C}$ have *effective unions*, i.e., the induced morphism $P \longrightarrow B$ from the pushout P of the pullback morphisms of a pair of (resp. pure) monos with codomain B is a (pure) mono. Typical examples are the categories of modules.

From [AR, 94] (see also 2.4 for the “pure” case), we know that in any locally finitely presentable category, there exists a cardinal γ such that:

- (a) each object C of $\mathcal{C}$ is a γ -filtered colimit $(u_i: C_i \longrightarrow C)$ with the C_i 's γ -presentable and the u_i 's monos (resp. pure monos), and
- (b) every subobject of a γ -presentable object has its domain γ -presentable.

Let S be the set of all (pure) subobjects of an object B , partially ordered by transfinite compositions of γ -presentable (pure) monos. Since S is closed under colimit of chains, any given (pure) mono $f: A \longrightarrow B$ must factorize $A \xrightarrow{t} C \xrightarrow{g} B$ through a maximal element $g: C \longrightarrow B$ of S , with t a transfinite composition of (pure) monos. We need to show that g is iso.

If not, then there exists a (pure) mono $b_0: B_0 \longrightarrow B$, with B_0 γ -presentable, which does not factorize through g (by (a)). We take the pullback (c_0, g_0) of (b_0, g) . All four morphisms are monos, and g_0 is γ -presentable because its domain is (see (b) above).

Then we take the pushout $(c'_0, g'_0: C \longrightarrow P)$ of (c_0, g_0) , and let $h: P \longrightarrow B$ be the induced morphism.

$$\begin{array}{ccc}
 C_0 & \xrightarrow{g_0} & B_0 \\
 \downarrow c_0 & & \downarrow b_0 \\
 & \nearrow c'_0 & \\
 C & \xrightarrow{g'_0} P & \xrightarrow{h} B \\
 & \searrow g & \\
 & &
 \end{array}$$

Then g'_0 is a γ -presentable, being the pushout of a γ -presentable morphism, and it is a (pure) mono, because g is. h is a (pure) mono, by the existence of effective unions of (pure) subobjects, hence g'_0 is an iso, by maximality of g . This is a contradiction because b_0 now factorizes through g .

REFERENCES

- [AHRT, 02] J. ADÁMEK, H. HERRLICH, J. ROSICKÝ, W. THOLEN, *On a generalized small-object argument for the injective subcategory problem*, Cahiers Top. Géom. Diff. Catég. 43 (2002), 83-106.
- [AHT, 96] J. ADÁMEK, H. HU, W. THOLEN, *On pure morphisms in accessible categories*, J. Pure Appl. Algebra 107 (1996), 1-8.
- [AR, 94] J. ADÁMEK, J. ROSICKÝ, *Locally Presentable and Accessible Categories*, Cambridge Univ. Press, Cambridge 1994.
- [Be, 67] A. BESSERRE, *Sur quelques propriétés des couples d'anneaux ou de modules*, J. Math. Pures et Appl. 46 (1967), 313-352.
- [Bk, 00] T. BEKE, *Sheafifiable homotopy model categories*, Math. Proc. Camb. Phil. Soc. 129 (2000), 447-475.
- [BR, 05] F. BORCEUX, J. ROSICKÝ, *Purity in algebra*, manuscript 2004.
- [Co, 79] M. COSTE, *Localisation, spectra and sheaf representation*, Appl. of Sheaves, Lect. Notes in Math. 753, Springer-Verlag, Berlin 1979, 212-238.
- [Ch, 05] B. CHORNY, *A generalization of Quillen's small object argument*, manuscript 2005, [http://arxiv.org \(math.AT/0401424\)](http://arxiv.org/math/AT/0401424).
- [CHK, 85] C. CASSIDY, M. HÉBERT, G.M. KELLY, *Reflective subcategories, localizations and factorization systems*, J. Austr. Math. Soc. 38 (Series A) (1985), 287-329.
- [Fa, 75] S. FAKIR, *Objets algébriquement clos et injectifs dans les catégories localement présentables*, Bull. Soc. Math. France, Mem. 42, 1975.
- [H, 98] M. HÉBERT, *Purity and injectivity in accessible categories*, J. Pure Appl. Algebra 129 (1998), 143-147.
- [H₁, 04] M. HÉBERT, *Algebraically closed and existentially closed substructures in categorical context*, Theory Appl. Categ. 12 (2004), 270-298.
- [H₂, 04] M. HÉBERT, *K-Purity and orthogonality*, Theory Appl. Categ. 12 (2004), 355-371.
- [HAR, 01] M. HÉBERT, J. ADÁMEK AND J. ROSICKÝ, *More on orthogonality in locally presentable categories*, Cahiers Top. Géom. Diff. Catég. 42 (2001), 51-80.
- [MP, 89] M. MAKKAÏ, R. PARÉ, *Accessible categories: the foundation of categorical model theory*, Cont. Math. 104, Amer. Math. Soc., Providence 1989.
- [NY, 03] W. NICHOLSON, M. YOUSIF, *Quasi-Frobenius Rings*, Cambridge Univ. Press, Cambridge 2003.
- [P, 03] G. PICAUVET, *Universally going-down rings, 1-split rings, and absolute integral closure*, Comm. Alg. 31 (2003), 4655-4685.
- [Po, 79] D. POPESCU, *Algebraically pure morphisms*, Rev. Roum. Math. Pures Appl., 24 (1979), 947-977.
- [Q, 67] D.G. QUILLEN, *Homotopical Algebra*, Lecture Notes in Math. 43, Springer-Verlag, Berlin 1967.
- [T, 01] W. THOLEN, *Essential weak factorization systems, Contributions to general algebra*, 13, 321-333, Heyn, Klagenfurt, 2001.